



# ANOGEN - A Division of YES Biotech Laboratories Ltd.

2355 Derry Road East, Unit 23, Mississauga, ON, Canada L5S 1V6 • Tel: (905) 677-9221 • Fax: (905) 677-0023

## Mouse Anti-Human IL-8 Monoclonal Antibody Datasheet

**Product Name:** mAb anti-Human IL-8

**Clone No.:** I8-61

**Catalogue No.:** MO-C40017E

**Quantity:** 0.5 mg/vial

**Description:** Mouse monoclonal antibody to human interleukin 8 (IL-8)

**Purification:** Protein G affinity purified

**Product Type:** Primary antibody

**Target Protein:** Human IL-8

**Immunogen:** Purified recombinant human IL-8

**Fusion Myeloma:** P3X63Ag8.653

**Specificity:** This antibody recognizes both recombinant and native human IL-8

**Species Reactivity:** Human, others not tested

**Cross-reactivity:** This antibody does not cross react with human Monocyte Chemotactic Activating Factor (MCAF) or RANTES (Regulated on Activation, Normal T-cell Expressed, and Secreted).

**Host / Isotype:** Mouse, IgG1 Kappa

**Formulation:** Lyophilized from a solution in 0.01M PBS, pH 7.0

**Reconstitution:** Double distilled water is recommended to adjust the final concentration to 1.00mg/mL

**Storage:** Store at -20°C

**Research Area:** Cytokine, interleukin and chemotaxis

**Background:** Interleukin 8 (IL-8), like IL-6, is secreted by macrophages and a variety of cells that express Toll-like receptors in response to the stimulation of pathogens. IL-8's primary function is to recruit neutrophils

and other target cells through chemotaxis to the infected site to eliminate pathogens. IL-8 causes increased intracellular Ca<sup>2+</sup>, release of reactive oxygen species, and other physiological changes required for migration and phagocytosis. IL-8 is also known to promote angiogenesis.


### Applications

**ELISA:** React with human IL-8.

**Neutralizing:** In chemotaxis assay, the antibody inhibited 97 % of the chemotactic activity of IL-8 on RB/293 cell using 5µg/ml of MAb and 10ng/ml of human IL-8 in assay. This antibody inhibited the chemotactic effect of human IL-8, and had no inhibitory effect to MIP 1-beta and RANTES in neutrophil chemotaxis assay, indicating that the neutralizing activity is specific. The antibody also neutralized IL-8 induced calcium influx in human granulocytes.

**Western Blot:** The antibody when used at 0.02-0.1mg/ml concentration, will allow visualization of 100ng/lane of recombinant human IL-8.

**Figure:** Western blot analysis of recombinant human IL-8 using clone I8-61.



### References:

If research is published using this product, please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product in the same category as gratitude.

**This product is for LABORATORY RESEARCH USE and further manufacture ONLY, and cannot be administered to human and animals for use in diagnostic and therapeutic procedures.**

**Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.**